

Hendry County Schools
Safe and Drug-Free Schools (SDFS)
Program

Contact Information

Larry Worth
P.O. Box 1980
LaBelle, FL 33975
(863) 674-4166 Ext 228
SunCom 744-4166
Fax (863) 674-4106
worthl@hendry.k12.fl.us

Key 1

NYA not yet available
LEP limited English proficient

District Data

7,593 public school students
13% LEP students
70% on free/reduced lunch
33% White, 16% Black,
48% Hispanic, 3% Other
11 schools in district (PreK-12th)

Project Goals

ATOD Prevention

By June 30, 2005, to decrease by 50% the percentage of students in grades 6-8 who report current use of alcohol (past 30 days), as measured by the Florida Youth Substance Abuse Survey (FYSAS). The baseline statistic from the 1999-2000 FYSAS was 20.4%.

By June 30, 2005, to decrease by 50% the percentage of students in grades 9-12 who report current use of alcohol (past 30 days), as measured by the Florida Youth Substance Abuse Survey (FYSAS). The baseline statistic from the 1999-2000 FYSAS was 43.8%.

By June 30, 2005, to decrease by 50% the percentage of students in grades 9-12 who

report binge drinking, as measured by the 2004-2005 FYSAS. The baseline statistic from the 1999-2000 FYSAS was 20.2%.

By June 30, 2005, to decrease by 50% the percentage of students in grades 6-8 who report binge drinking, as measured by the 2004-2005 FYSAS. The baseline statistic from the 1999-2000 FYSAS was 13.8%.

By June 30, 2005, to increase by 2.5 years the mean age of first use of alcohol in students in grades 6-8, as measured by the 2004-2005 FYSAS. The baseline statistic from the 1999-2000 FYSAS was 11 years of age.

By June 30, 2005, to increase by 2.5 years the mean age of first use of alcohol in students in grades 9-12, as measured by the 2004-2005 FYSAS. The baseline statistic from the 1999-2000 FYSAS was 12.9 years of age.

By June 30, 2005, to decrease by 50% the percentage of students in grades 6-8 who report current use of cigarettes (past 30 days), as measured by the 2004-2005 FYSAS. The baseline statistic from the 1999-2000 FYSAS was 9.4%.

By June 30, 2005, to decrease by 50% the percentage of students in grades 9-12 who report current use of cigarettes (past 30 days), as measured by the 2004-2005 FYSAS. The baseline statistic from the 1999-2000 FYSAS was 21.3%.

By June 30, 2005, to decrease by 50% the percentage of students in grades 6-8 who report current use of smokeless tobacco (past 30 days), as measured by the 2004-2005 FYSAS. The baseline statistic from the 1999-2000 FYSAS was 6.8%.

By June 30, 2005, to decrease by 50% the percentage of students in grades 9-12 who report current use of smokeless tobacco (past 30 days), as measured by the 2004-2005 FYSAS. The baseline statistic from the 1999-2000 FYSAS was 5.9%.

Safe and Drug Free Schools 2004-2005 District Profiles

By June 30, 2005, to increase by 2.5 years the mean age of first use of tobacco in students in grades 6-8, as measured by the 2004-2005 FYSAS. The baseline statistic from the 1999-2000 FYSAS was 11 years of age.

By June 30, 2005, to increase by 2.5 years the mean age of first use of tobacco in students in grades 6-12, as measured by the 2004-2005 FYSAS. The baseline statistic from the 1999-2000 FYSAS was 12.5 years of age.

By June 30, 2005, to decrease by 50% the percentage of students in grades 6-8 who report current use of marijuana (past 30 days), as measured by the 2004-2005 FYSAS. The baseline statistic from the 1999-2000 FYSAS was 4.9%.

By June 30, 2005, to decrease by 50% the percentage of students in grades 9-12 who report current use of marijuana (past 30 days), as measured by the 2004-2005 FYSAS. The baseline statistic from the 1999-2000 FYSAS was 17.7%.

By June 30, 2005, to increase by 2.5 years the mean age of first use of marijuana in students in grades 6-8, as measured by the 2004-2005 FYSAS. The baseline statistic from the 1999-2000 FYSAS was 11.9 years of age.

By June 30, 2005, to increase by 2.5 years the mean age of first use of marijuana in students in grades 9-12, as measured by the 2004-2005 FYSAS. The baseline statistic from the 1999-2000 FYSAS was 13.6 years of age.

By June 30, 2005, to decrease by 50% the percentage of students in grades 6-8 who report current use of any illicit drug excluding marijuana (past 30 days), as measured by the 2004-2005 FYSAS. The baseline statistic from the 1999-2000 FYSAS was 7.6%.

By June 30, 2005, to decrease by 50% the percentage of students in grades 9-12 who report current use of any illicit drug excluding marijuana (past 30 days), as measured by the 2004-2005 FYSAS. The baseline statistic from the 1999-2000 FYSAS was 12.3%.

Violence Prevention

By June 30, 2005, to decrease by 20% the number of fighting incidents per 1000 for students in grades PK-12 as measured the 2004-2005 School Environmental Safety Incident Report (SESIR) data. The 1999-2000 baseline statistic for the number of fighting incidents per 1000 for students in grades PK-12 in Hendry County was 18.59.

Key 2

- √ Proven Program
- ◆ Promising Program
- AP** ATOD Prevention
- VP** Violence Prevention
- PI** Parent Involvement

SDFS Programs

- Too Good for Drugs, √, **AP, PI**
- Too Good for Violence, ◆, **VP, PI**
- Too Good for Drugs and Violence-High School, ◆, **AP, VP, PI**
- Teen Summit, ◆, **AP, VP**

SDFS Collaborations

- Hendry County SDFS Coalition-Community Awareness, **PI**
- Hendry County Sheriff's Department Students Taking a New Direction (STAND), **AP, VP, PI**
- Hendry County Sheriff's Department School Resource Officers, **AP, VP, PI**
- U.S. Department of Education-Heartland Consortium, **AP, VP, PI**