

# ARBITRAGE-FREE OPTION PRICING MODELS

DENIS BELL<sup>1</sup>and SCOTT STELLJES

(December 26, 2008)

## Abstract

We describe a scheme for constructing explicitly solvable arbitrage-free models for stock price. This is used to study a model similar to one introduced by Cox and Ross, where the volatility of the stock is proportional to the square root of the stock price. We derive a formula for the value of a European call option based on this model and give a procedure for estimating parameters and for testing the validity of the model.

*Keywords and phrases:* Arbitrage-free, Black–Scholes formula, European call option.

## 1. Introduction

The famous Black–Scholes model [1] for option pricing is based on the assumption that the value  $S$  of a stock follows an Itô equation of the form

$$dS = \mu S dt + \sigma S dw. \quad (1)$$

Here  $w$  is a standard Wiener process and  $\mu$  and  $\sigma$  are two parameters representing, respectively, the drift and the volatility of the stock. This leads to the well-known Black–Scholes formula for determining the value  $V$  of a European call option, i.e. the right to purchase the stock at a price  $k$  at a future time  $T$ ,

$$V = S_0 \Phi\left(\frac{\log(S_0/k) + (r + \sigma^2/2)T}{\sigma\sqrt{T}}\right) - ke^{-rT} \Phi\left(\frac{\log(S_0/k) + (r - \sigma^2/2)T}{\sigma\sqrt{T}}\right). \quad (2)$$

In this formula,  $S_0$  denotes the present value of the stock,  $\Phi$  is the cumulative distribution function of the standard Gaussian distribution and  $r$  is the risk-free interest rate.

---

<sup>1</sup>RESEARCH PARTIALLY SUPPORTED BY NSF GRANT DMS-0451194.

An essential feature of option pricing is the calculation of expectation with respect to an underlying probability measure with respect to which the model (1) is *arbitrage-free*. This is the condition that the process  $e^{-rt}S_t$  is a *martingale*. In particular, the expected value of  $S_t$  at any time  $t$  is precisely the future value at time  $t$  of a risk-free bond with present value  $S_0$ , i.e.

$$E[S_t] = S_0e^{rt}. \quad (3)$$

The arbitrage-free condition effectively forces the replacement of  $\mu$  in equation (1) by  $r$  in the subsequent computations of expectations.

Alternatives to the Black–Scholes model (1) have been proposed. In particular Cox and Ross [2] introduced the model

$$dS = \mu Sdt + \sigma\sqrt{S}dw. \quad (4)$$

They obtained this equation as the diffusion limit of a sequence of jump processes with intensity tending to infinity. In contrast to (1), equation (4) is not solvable in closed form by elementary means. Thus there is no direct analogue of the Black–Scholes formula (2) for the Cox–Ross model.

In Section 2 we describe a method for constructing a class of *solvable* arbitrage-free models for stock price. Our starting point is the following stochastic Bernoulli equation of *Stratonovich* type

$$d\tilde{S} = \mu\tilde{S}dt + \sigma\tilde{S}^p \circ dw, \quad 1/2 \leq p \leq 1. \quad (5)$$

In view of the fact that Stratonovich differentials transform in the same way as classical differentials, equation (5) can be solved explicitly by elementary methods (cf. the Zvonkin, Doss–Sussmann method, Karatzas and Shreve [5, Proposition 2.21], Rogers and Williams [7, Theorem 28.2]). This is done in Theorem 1.

The process  $\tilde{S}$  will not generally satisfy the arbitrage-free condition. In Theorem 2, we construct a function  $G$  such that the process  $S_t \equiv G(\tilde{S}_t, t)$  *does* satisfy this condition. This yields a second-order partial differential equation for  $G$  that is similar to the classical Black–Scholes equation.

In Section 3 we consider the extreme values  $p = 1$  and  $p = 1/2$ , which are particularly tractable to this analysis. In the linear case  $p = 1$  our approach is shown to yield the Black–Scholes model. We then focus on the case  $p = 1/2$  studied by Cox and Ross. In this case the partial differential equation for  $G$  turns out to have an especially simple form and we are able to solve it explicitly. This results in a formula (equation (17)) for the value of a European call option analogous to the Black–Scholes formula (2).

Finally, in Section 4, we give a method for estimating the volatility parameter  $\sigma$  from a set of data and for testing the validity of the model.

It should be pointed out that the idea of using Stratonovich calculus in mathematical finance is not new. A main objective of this paper is to use

this methodology to provide a unified treatment of the Black–Scholes and Cox–Ross models that is self-contained and, at the same time, elementary enough to be comprehensible to a wide variety of readers.

## 2. Arbitrage-free models

Throughout this section, let  $p = m/n$  denote a rational in the interval  $[1/2, 1]$ , with  $m$  odd and  $n$  even. We introduce the following stochastic differential equation as a tentative model for stock price

$$d\tilde{S} = r\tilde{S}dt + \sigma\tilde{S}^p \circ dw, \quad (6)$$

where  $\circ dw$  denotes the Stratonovich differential. The following relationship exists between Itô and Stratonovich differentials (cf. Klebaner [6, Theorem 5.20]): for a random process  $\xi$  with a stochastic differential  $d\xi = adw + bdt$ , where  $a$  and  $b$  are continuous adapted processes, we have

$$\begin{aligned} \xi \circ dw &= \xi dw + \frac{1}{2}d[\xi, w] \\ &= \xi dw + \frac{1}{2}adt \end{aligned} \quad (7)$$

where the bracket  $[\xi, w]$  denotes the quadratic covariation of the semimartingales  $\xi$  and  $w$ . It follows from this and Itô's formula (cf. e.g. [6, Theorem 4.16]) that Stratonovich differentials transform under composition with smooth maps in the same way as classical differentials, i.e. by the standard chain rule. Thus equation (6) may formally be regarded as an ordinary differential equation in  $\tilde{S}$ . As such, it is solvable by elementary differential equation methods, i.e. variation of parameters and separation of variables. This gives the following result.

THEOREM 1. *Define*

$$\tilde{S}_t = e^{rt} \left\{ (1-p)\sigma \int_0^t e^{r(p-1)u} dw_u + \tilde{S}_0^{1-p} \right\}^{\frac{1}{1-p}}, \quad t \geq 0. \quad (8)$$

Then  $\tilde{S}_t$  is the solution to equation (6).

*Proof.* Note that, although the quantity inside the rational power  $1/(1-p)$  in (8) may be negative, the assumption on  $p$  ensures that  $\tilde{S}_t$  is real and non negative.

We verify that (8) is the solution to equation (6) as follows. Write

$$E_t \equiv \left\{ (1-p)\sigma \int_0^t e^{r(p-1)u} dw_u + \tilde{S}_0^{1-p} \right\}^{\frac{1}{1-p}}$$

and note that the function  $f(x) = x^{\frac{1}{1-p}}$  is  $C^2$  on  $\mathbf{R}$ . Applying Itô's formula to compute the stochastic differential of the process  $\tilde{S}_t = e^{rt}E_t$  and using (7) gives

$$\begin{aligned} d\tilde{S} &= re^{rt}E_t dt + \sigma e^{rpt}E_t^p dw + \frac{p\sigma^2}{2}e^{rt(2p-1)}E_t^{2p-1}dt \\ &= (r\tilde{S} + \frac{p\sigma^2}{2}\tilde{S}^{2p-1})dt + \sigma\tilde{S}^p dw \\ &= r\tilde{S}dt + \sigma\tilde{S}^p dw + \frac{1}{2}d[\sigma\tilde{S}^p, w] \\ &= r\tilde{S}dt + \sigma\tilde{S}^p \circ dw \end{aligned}$$

as required.

As remarked in Section 1, the process  $\tilde{S}_t$  will not generally satisfy the arbitrage-free condition and hence is not a feasible model for stock price. We therefore seek a function  $G$  such that  $S_t \equiv G(\tilde{S}_t, t)$  is arbitrage-free.

**THEOREM 2.** *Suppose that  $G : \mathbf{R} \times \mathbf{R} \mapsto \mathbf{R}$  is such that  $G(s, t)$  is  $C^2$  in  $s$ ,  $C^1$  in  $t$ , and satisfies the partial differential equation*

$$G_t + \left( rs + \frac{p\sigma^2 s^{2p-1}}{2} \right) G_s + \frac{\sigma^2 s^{2p}}{2} G_{ss} = rG. \quad (9)$$

Suppose further that there exists  $n$  such that for every  $T > 0$

$$\sup_{0 \leq t \leq T} |G_s(s, t)| \leq C|s|^n, \quad \forall s \in \mathbf{R} \quad (10)$$

where  $C$  is a constant depending only on  $T$ .

Let  $S_t = G(\tilde{S}_t, t)$ . Then the process  $e^{-rt}S_t$  is a martingale.

*Proof.* We have

$$\begin{aligned} d(e^{-rt}S_t) &= e^{-rt} \left( (G_t(\tilde{S}_t, t) - rG(\tilde{S}_t, t))dt + G_s(\tilde{S}_t, t)d\tilde{S}_t \right) \\ &= e^{-rt} \left( (G_t(\tilde{S}_t, t) - rG(\tilde{S}_t, t))dt + G_s(\tilde{S}_t, t)(r\tilde{S}dt + \sigma\tilde{S}^p \circ dw) \right). \quad (11) \end{aligned}$$

Applying (7) to convert equation (11) to Itô form then using (9), we obtain

$$\begin{aligned} d(e^{-rt}S_t) &= e^{-rt} \left[ G_t(\tilde{S}_t, t) + \left( r\tilde{S}_t + \frac{p\sigma^2}{2}\tilde{S}_t^{2p-1} \right) G_s(\tilde{S}_t, t) \right. \\ &\quad \left. + \frac{\sigma^2 \tilde{S}_t^{2p}}{2} G_{ss}(\tilde{S}_t, t) - rG(\tilde{S}_t, t) \right] dt + e^{-rt} G_s(\tilde{S}_t, t) \sigma \tilde{S}_t^p dw \\ &= e^{-rt} G_s(\tilde{S}_t, t) \sigma \tilde{S}_t^p dw. \end{aligned}$$

Thus

$$e^{-rt}S_t = S_0 + \sigma \int_0^t e^{-ru} G_s(\tilde{S}_u, u) \tilde{S}_u^p dw. \quad (12)$$

Now the indefinite Itô integral  $\int_0^s f dw$ ,  $s \leq t$  is a martingale (cf. [6, Page 100]) provided  $f$  satisfies

$$\int_0^t E[f(u)^2] du < \infty. \quad (13)$$

Using the estimate (cf. Gikhman and Skorohod [4])

$$E\left[\sup_{0 \leq s \leq T} \left| \int_0^s f(u) dw \right|^{2m}\right] \leq C_m \int_0^T E[|f(u)|^{2m}] du$$

for some constant  $C_m$ , and condition (10), it is easy to check that the integrand in (12) satisfies (13). Thus the result holds.

### 3. Examples

There are two values,  $p = 1$  and  $p = 1/2$ , where equation (9) is solvable in closed form. Firstly, in the case  $p = 1$ , (9) reduces to

$$G_t + \left(r + \frac{\sigma^2}{2}\right) s G_s + \frac{\sigma^2 s^2}{2} G_{ss} = rG.$$

It is easy to check that

$$G(s, t) = se^{-\sigma^2 t/2}$$

is a solution to this equation and this function  $G$  clearly satisfies condition (10). Solving equation (6) in the case  $p = 1$  yields

$$\tilde{S}_t = S_0 \exp(rt + \sigma w_t).$$

Thus

$$S_t = G(\tilde{S}_t, t) = S_0 \exp\left(\left(r - \frac{\sigma^2}{2}\right)t + \sigma w_t\right).$$

We also note that when  $p = 1$ , the Itô equation for  $S$  is

$$dS = rSdt + \sigma Sdw$$

and the equation that gives rise to this, prior to the imposing of the arbitrage-free condition, is precisely (1). Thus both our model and its solution reduce to the classical Black–Scholes theory in the case  $p = 1$ .

We now turn to the case  $p = 1/2$  studied by Cox and Ross (this will be referred to in the sequel as the *fractional* model). In this case, equation (9) becomes

$$G_t + \left(rs + \frac{\sigma^2}{4}\right) G_s + \frac{\sigma^2 s}{2} G_{ss} = rG$$

which has the simple solution

$$G(s, t) = s + \frac{\sigma^2}{4r}. \quad (14)$$

Again,  $G$  satisfies (10).

Assume that  $S_0 > \sigma^2/4r$ . Combining (14) and (8) with  $p = 1/2$  gives

$$S_t = G(\tilde{S}_t, t) = e^{rt} \left\{ \frac{\sigma}{2} \int_0^t e^{-ru/2} dw_u + \sqrt{S_0 - \frac{\sigma^2}{4r}} \right\}^2 + \frac{\sigma^2}{4r} \quad (15)$$

The Itô equation for  $S$  is

$$dS = rSdt + \sigma \sqrt{S - \frac{\sigma^2}{4r}} dw.$$

A natural candidate for the original model of stock price that gives rise to this formula after the imposing of the arbitrage-free condition, is

$$dS = \mu Sdt + \sigma \sqrt{S - \frac{\sigma^2}{4r}} dw. \quad (16)$$

Formula (15) yields a value  $V$  for a call option with exercise price  $k$  at future time  $T$ . The Itô integral in (15) is a Gaussian random variable with zero mean and variance  $(1 - e^{-rt})/r$ . An elementary calculation shows that

$$V = e^{-rT} E[\max(S_T - k, 0)]$$

is given by

$$V = \frac{v}{2\pi} \left( e^{-d_1^2/2} (vd_1 + 2a) - e^{-d_2^2/2} (vd_2 + 2a) \right) + \left( (v^2 + a^2) + e^{-rT} (b - k) \right) \left( 1 - \Phi(d_1) + \Phi(d_2) \right) \quad (17)$$

where  $\Phi$  is the cumulative distribution function of the standard normal distribution,

$$a = \sqrt{S_0 - \frac{\sigma^2}{4r}}, \quad b = \frac{\sigma^2}{4r}, \quad v = \frac{\sigma}{2} \sqrt{\frac{1 - e^{-rT}}{r}},$$

$$d_1 = \frac{e^{-rT/2} \sqrt{k - b} - a}{v}, \quad d_2 = \frac{-e^{-rT/2} \sqrt{k - b} - a}{v}.$$

#### 4. Parameter estimation and model testing

In order to use formula (17), a mechanism is required to estimate the volatility  $\sigma$  for a given stock from a set of data. We propose the following scheme, based on a list of opening prices  $S_t$  for the stock on successive days  $t = 1, \dots, N$ . Since the data set is discrete, it is natural to approximate equation (16) by the *difference* equation

$$\Delta_t S = \mu S_t + \sigma \sqrt{S_t - \frac{\sigma^2}{4r}} \Delta_t w$$

where  $\Delta_t S$  and  $\Delta_t w$  denote  $S_{t+1} - S_t$  and  $w_{t+1} - w_t$ , respectively. Solving for  $\Delta_t w$ , we see that, if the model is valid, then the quantities

$$Z_t \equiv \frac{\Delta_t S - \mu S_t}{\sigma \sqrt{S_t - \frac{\sigma^2}{4r}}} \quad (18)$$

form an (approximately) independent set of standard Gaussian random variables. Equating the mean to zero and the variance to 1 of the sample values  $Z_t, t = 1, \dots, N$  gives the following equations in  $\mu$  and  $\sigma$

$$\sum_{t=1}^N Z_t = 0$$

$$\sum_{t=1}^N Z_t^2 = N - 1.$$

These equations can be solved numerically to yield estimates  $\hat{\mu}$  and  $\hat{\sigma}$  of the parameters  $\mu$  and  $\sigma$  and the estimated value  $\hat{\sigma}$  can then be used in place of  $\sigma$  in (17) to compute the option value  $V$ . Furthermore, the validity of the fractional model can be tested by applying a standard normality test, e.g. Shapiro–Wilk to the quantities  $Z_t$  in (18), again using the estimated values for  $\mu$  and  $\sigma$  in place of the actual values.

In conclusion, we note that Delbaen and Shirakawa [3] have recently used a Bessel process to study the law of the process  $\hat{S}_t$  defined by the *Itô* equation

$$d\hat{S} = r d\hat{S} dt + \sigma \hat{S}^p dw.$$

Here  $p$  is assumed to lie in the range  $(0, 1)$ . The analysis in [3] results in an expression for the law of the random variable  $\hat{S}_t, t > 0$  as an infinite series. This result is then used to obtain a formula [3, Equation 3.21] for the price of a European call option based on the stock price  $\hat{S}_t$ , as the difference of two infinite series. The relationship between the option pricing formula in [3] and its counterpart (equation (17)) in the present work is unclear at this time. This issue will be studied in a later paper.

## References

- [1] F. Black and M. J. Scholes, The pricing of options and corporate liabilities, *Journal of Political Economy* 81, no. 3 (1973), 635-654.
- [2] J. C. Cox and S. A. Ross, The valuation of options for alternative stochastic processes, *Journal of Financial Economics* 3 (1976), 145-166.
- [3] F. Delbaen and H. Shirakawa, A note of option pricing for constant elasticity of variance model, preprint No. 96-03.
- [4] I. I. Gikhman and A. V. Skorohod, *Stochastic Differential Equations*, Springer-Verlag, Berlin, 1972.
- [5] I. Karatzas and S. Shreve, *Methods of Mathematical Finance*, Springer-Verlag, New York, 1998.
- [6] F. Klebaner, *Introduction to Stochastic Calculus with Applications*. Second edition. Imperial College Press, London, 2005.
- [7] L. C. G. Rogers and D. Williams, *Diffusions, Markov Processes, and Martingales*, Vol. 2, Wiley, New York, 1987.

Department of Mathematics and Statistics  
University of North Florida  
1 UNF Drive  
Jacksonville, FL 32224  
U. S. A.